### **Environmental Protection Agency**

Cattle, fat Cattle, kidney Cattle, meat Cattle, meat byproducts Goat, fat Goat, kidney Goat, meat Goat, meat Hog, kidney Hog, meat	arts per million
Horse, kidney Horse, meat byproducts Milk Sheep, fat Sheep, kidney	0.1 0.5 0.1 0.1 0.1 0.1 0.1 0.1 0.1 0.1 0.1 0.1
Sheep, meat byproducts	0.1 0.1

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[64 FR 70191, Dec. 16, 1999, as amended at 66 FR 64773, Dec. 14, 2001; 67 FR 51097, Aug. 7, 2002]

# § 180.429 Chlorimuron ethyl; tolerances for residues.

(a) General. Tolerances are established for residues of the herbicide chlorimuron ethyl, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified in the following table is to be determined by measuring only chlorimuron ethyl, ethyl 2-[[[(4-chloro-6-methoxypyrimidin-2yl)amino]carbonyl]sulfonyl]benzoate]

2yl)amino]carbonyl]sulfonyl]benzoate] in or on the following commodities:

Commodity	Parts per million
Berry, low growing, except strawberry, subgroup 13–07H	0.02
Corn, field, forage	0.02
Corn, field, grain	0.01
Corn, field, stover	2.0
Grain, aspirated fractions	3.0
Peanut	0.02
Soybean, forage	0.45
Soybean, hay	1.8
Soybean, seed	0.05

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]

(d) Indirect or inadvertent residues. [Reserved]

[74 FR 10494, Mar. 11, 2009, as amended at 74 FR 67087, Dec. 18, 2009]

## §180.430 Fenoxaprop-ethyl; tolerances for residues.

(a) General. Tolerances are established for residues of the herbicide fenoxaprop-ethyl, including its metabolites and degradates, in or on the commodities in the table in this paragraph. Compliance with the tolerance levels specified in this paragraph is to be determined by measuring only the sum of fenoxaprop-ethyl, (±)-ethyl 2-[4-[(6-chloro-2-

benzoxazolyl)oxy]phenoxy]propanoate, and its metabolites, 2-[4-[(6-chloro-2-benzoxazolyl)oxy]phenoxy]propanoic acid and 6-chloro-2,3-dihydrobenzoxazol-2-one, calculated as

dihydrobenzoxazol-2-one, calculated as the stoichiometric equivalent of fenoxaprop-ethyl, in or on the commodity.

Commodity	Parts per million
Barley, grain	0.05
Barley, straw	0.1
Cattle, fat	0.05
Cattle, meat byproducts	0.05
Cattle, meat	0.05
Cotton, undelinted seed	0.05
Goat, fat	0.05
Goat, meat byproducts	0.05
Goat, meat	0.05
Hog, fat	0.05
Hog, meat byproducts	0.05
Hog, meat	0.05
Horse, fat	0.05
Horse, meat byproducts	0.05
Horse, meat	0.05
Milk	0.02
Peanut	0.05
Peanut, hulls	0.05
Rice, grain	0.05
Sheep, fat	0.05
Sheep, meat byproducts	0.05
Sheep, meat	0.05
Soybean	0.05
Wheat, grain	0.05
Wheat, straw	0.50

(b) Section 18 emergency exemptions. Time-limited tolerances are established for residues of the herbicide fenoxaprop-ethyl, including its metabolites and degradates, in or on the commodities in the table in this paragraph in connection with use of fenoxaprop-ethyl under section 18 emergency exemptions granted by EPA. Compliance with the tolerance levels specified in this paragraph is to

#### § 180.431

be determined by measuring only the sum of fenoxaprop-ethyl, (±)-ethyl 2-[4-[(6-chloro-2-

benzoxazolyl)oxy]phenoxy]propanoate, and its metabolites, 2-[4-[(6-chloro-2-benzoxazolyl)oxy]phenoxy]propanoic acid and 6-chloro-2,3-dihydrobenzoxazol-2-one, calculated as the stoichiometric equivalent of fenoxaprop-ethyl, in or on the commodity. The tolerances expire and are revoked on the dates specified in the table in this paragraph.

Commodity	Parts per million	Expiration/ revocation date
Grass, forage	0.05 0.05	12/31/16 12/31/16

(c) Tolerances with regional registrations. Tolerances with regional registration, as defined in §180.1(1), are established for residues of the herbicide fenoxaprop-ethyl, including its metabolites and degradates, in or on the commodities in the table in this paragraph when fenoxaprop-ethyl is used in the states of Oregon, Washington, and Utah. Compliance with the tolerance levels specified in this paragraph is to be determined by measuring only the sum of fenoxaprop-ethyl, (±)-ethyl 2-[4-[(6-chloro-2-

benzoxazolyl)oxy]phenoxy]propanoate, and its metabolites, 2-[4-[(6-chloro-2-benzoxazolyl)oxy]phenoxy]propanoic acid and 6-chloro-2,3-dihydrobenzoxazol-2-one, calculated as the stoichiometric equivalent of fenoxaprop-ethyl, in or on the commodity

Commodity	Parts per million
Grass, hay	0.09

(d) Indirect or inadvertent residues. [Reserved]

[63 FR 1377, Jan. 9, 1998, as amended at 63 FR 19837, Apr. 22, 1998; 73 FR 33718, June 13, 2008; 75 FR 80346, Dec. 22, 2010; 76 FR 23495, Apr. 27, 2011; 78 FR 78748, Dec. 27, 2013; 79 FR 26164, May 7, 2014]

## § 180.431 Clopyralid; tolerances for residues.

(a) General. Tolerances are established for residues of the herbicide clopyralid, including its metabolites

and degradates, in or on the commodities in the table below from its application in the acid form or in the form of its salts. Compliance with the tolerance levels specified below is to be determined by measuring only clopyralid, (3,6-dichloro-2-pyridinecarboxylic acid), in or on the following commodities:

Commodity	Parts per million
Apple	0.05
Asparagus	1.0
Barley, bran	12
Barley, grain	3.0 9.0
Barley, hay Barley, pearled barley	12
Barley, straw	9.0
Beet, garden, tops	3.0
Beet, garden, roots	4.0
Beet, sugar, molasses	10
Beet, sugar, roots	2.0
Beet, sugar, tops	3.0
Brassica, head and stem, subgroup 5A Brassica, leafy greens, subgroup 5B	2.0 5.0
Bushberry subgroup 13-07B	0.50
Canola, meal	6.0
Canola, seed	3.0
Cattle, fat	1.0
Cattle, liver	3.0
Cattle, meat	1.0
Cattle, meat byproducts, except liver  Corn, field, forage	36.0 3.0
Corn, field, grain	1.0
Corn, field, milled byproducts	1.5
Corn, field, stover	10.0
Corn, pop, grain	1.0
Corn, pop, stover	10.0
Corn, sweet, forage	7.0
Corn, sweet, kernel plus cob with husks re- moved	1.0
Corn, sweet, stover	10.0
Cranberry	4.0
Egg	0.1
Flax, meal	6.0
Fruit, stone, group 12	0.5
Goat, fat	1.0 3.0
Goat, meat	1.0
Goat, meat byproducts, except liver	36.0
Grass, forage	500.0
Grass, hay	500.0
Hog, fat	0.2
Hog, meat	0.2
Hog, meat byproducts	0.2 5.0
Horse, fat	1.0
Horse, liver	3.0
Horse, meat	1.0
Horse, meat byproducts, except liver	36.0
Milk	0.2
Oat, forage Oat, grain	9.0 3.0
Oat, greats/rolled oats	12
Oat, straw	9.0
Peppermint, tops	3.0
Plum, prune, dried	1.5
Poultry, fat	0.2
Poultry, meat	0.2
Poultry, meat byproducts	0.2 3.0
Rapeseed, meal	6.0
Rapeseed, subgroup 20A, except gold of pleas-	0.0
ure	3.0